

Core 2 Indices & Logarithms Answers

3(a)	$\log 0.8^x = \log 0.05$ $x \log_{10} 0.8 = \log_{10} 0.05$ oe $x = 13.425$ to 3dp	$x = \log_{0.8} 0.05$ (M1) 13.425 (A2) (else A1 for 1 or 2dp)	M1 A1 A1	3	NMS: SC B2 for 13.425 or better (B1 for 13.4 or 13.43; 13.42) Condone greater accuracy
(b)(i)	$\frac{a}{1-r}$ $\frac{a}{1-r} = 5a \Rightarrow a = 5a(1-r)$ $\Rightarrow 1 = 5(1-r) \Rightarrow r = \frac{4}{5} = 0.8$	M1 A1 A1	3	$S_{\infty} = \frac{a}{1-r}$ <u>used</u> Or better AG (be convinced)	
(ii)	$n^{\text{th}} \text{ term} = 20 \times (0.8)^{n-1}$ $n^{\text{th}} \text{ term} < 1 \Rightarrow 0.8^{n-1} < \frac{1}{20}$ oe Least n is 15	M1 A1 A1F	3	Condone $20 \times (0.8)^n$. $0.8^{n-1} < 0.05$ or $0.8^{n-1} = k$, where $k=0.05$ or k rounds up to 0.050 If not 15, ft on integer part of [answer (a)+2] provided $n > 2$ SC 3/3 for 15 if no error SC $n^{\text{th}} \text{ term} = 16^{n-1}$ M1A0A0	
Total				9	

7(a)	$2 \log_a n - \log_a (5n - 24) = \log_a 4$ $\Rightarrow \log_a n^2 - \log_a (5n - 24) = \log_a 4$ $\Rightarrow \log_a \left[\frac{n^2}{5n - 24} \right] = \log_a 4$ $\Rightarrow \frac{n^2}{5n - 24} = 4$ $\Rightarrow n^2 - 20n + 96 = 0$	M1 M1 A1	3	A law of logs used A second law of logs used leading to both sides being single log terms or single log term on LHS with RHS=0 CSO. AG
(b)	$\Rightarrow (n - 8)(n - 12) = 0$ $\Rightarrow n = 8, 12$	M1 A1	2	Accept alternatives eg formula, completing of sq..
Total			5	

5(a)	$\log_a x = \log_a 6^2 - \log_a 3$	M1		One law of logs used correctly
	$\log_a x = \log_a \left(\frac{6^2}{3}\right)$	M1		A second law of logs used correctly
	$\log_a x = \log_a \frac{36}{3} \Rightarrow x = 12$	A1	3	CSO AG
(b)	$\log_a y + \log_a 5 = 7 \Rightarrow \log_a 5y = 7$	M1		
	$\Rightarrow 5y = a^7$ or $y = \frac{1}{5}a^7$ or $a = (5y)^{1/7}$	m1 A1	3	Eliminates logs Accept these forms
Total			6	

3(a)(i)	$\{p\} = 2$	B1		Condone '64=8 ² '
(ii)	$\{q\} = -2$	B1ft		Ft on '-p' if q not correct
(iii)	$\{r\} = 0.5$	B1	3	Condone ' $\sqrt{8} = 8^{0.5}$ '
(b)	$\frac{8^x}{8^{0.5}} = 8^{-2} \Rightarrow 8^{x-0.5} = 8^{-2}$ OE	M1		Using parts (a) & valid index law to stage $8^c = 8^d$ (PI)
	$\Rightarrow x - 0.5 = -2 \Rightarrow x = -1.5$	A1ft	2	Ft on c's (q+r) if not correct (Accept correct answer without working)
ALT: $\log 8^x = \log k, x \log 8 = \log k; x = -1.5$				(M1 A1)
Total			5	

1(a)(i)	x^2	B1	1	
(ii)	$x^{\frac{1}{2}} = \sqrt{x}$	B1	1	Accept either form
(iii)	x^3	B1	1	
(b)(i)	$\int 3x^{\frac{1}{2}} dx = \frac{3}{\frac{3}{2}} x^{\frac{3}{2}} (+c)$	M1 A1		Index raised by 1 Simplification not yet required
	$= 2x^{\frac{3}{2}} + c$	A1	3	Need simplification <u>and</u> the + c OE
(ii)	$\int_1^9 3x^{\frac{1}{2}} dx = (2 \times 9^{\frac{3}{2}}) - (2 \times 1^{\frac{3}{2}})$	M1		F(9) - F(1), where F(x) is candidate's answer to (b)(i) [or clear recovery]
	$= 52$	A1ft	2	Ft on (b)(i) answer of form $kx^{1.5}$ i.e. $26k$
Total			8	

8(a)	$\log_a n = \log_a 3(2n-1)$ $\Rightarrow n = 3(2n-1)$ $\Rightarrow 3 = 5n \Rightarrow n = \frac{3}{5}$	M1 m1		OE Log law used PI by next line OE, but must not have any logs.
(b)(i)	$\log_a x = 3 \Rightarrow x = a^3$	A1	3	
(ii)	$\log_a y - \log_a 2^3 = 4$	B1	1	
	$\log_a \frac{y}{2^3} = 4$ $\begin{cases} xy = a^7 \times a^{(3\log_a 2)} \\ \text{or} \\ y = a^4 \times a^{(3\log_a 2)} \end{cases}$	M1		$3\log 2 = \log 2^3$ seen or used any time in (ii) Correct method leading to an equation involving y (or xy) and a log but not involving + or -
	$\frac{y}{2^3} = a^4$ $\begin{cases} xy = a^7 \times 2^3 \\ \text{or} \\ y = a^4 \times 2^3 \end{cases}$	m1		Correct method to eliminate ALL logs e.g. using $\log_a N = k \Rightarrow N = a^k$ or using $a^{\log_a c} = c$
	$by = a^3 \times 8a^4$ or $8a^7$	A1	4	
	Total		8	