

Core 2 Sequences Answers

5(a) $150 = 200p + q$ $120 = 150p + q$ $p = 0.6$ $q = 30$	M1 A1 m1 A1 B1	5	Either equation Both (condone embedded values for the M1A1) Valid method to solve two simultaneous eqns in p <u>and</u> q to find either p or q AG (condone if left as a fraction)
(b) $u_4 = 102$	B1F \checkmark	1	Ft on $(72 + q)$
(c) $L = pL + q ; L = 0.6L + 30$ $L = \frac{q}{1-p}$ $L = 75$	M1 m1 A1F \checkmark	3	Ft on $2.5q$
Total		9	

3(a) (Tenth term) = $a + (10-1)d$ = $1 + 9(6) = 55$	M1 A1	2	NMS or rep. addn. B2 CAO SC if M0 award B1 for $6n-5$ OE
(b)(i) $S_n = \frac{n}{2}[2 + (n-1)d]$ $\frac{n}{2}[2 + 6n - 6] = 7400$	M1 A1		Formula for $\{S_n\}$ with either $a = 1$ or $d = 6$ substituted Eqn formed with some expansion of brackets
$3n^2 - 2n = 7400 \Rightarrow 3n^2 - 2n - 7400 = 0$	A1	3	CSO AG
(ii) $(3n+148)(n-50) = 0$ $\Rightarrow n = 50$	M1 A1	2	Formula/factorisation OE NMS single ans. 50.. B2 CAO NMS 50 and $-49.3(3\dots)$ B1 CAO
Total		7	

4(a)	$(1-2x)^4 = (1)^4 + 4(1)^3(-2x) + 6(1^2)(-2x)^2 + [4(1)(-2x)^3 + (-2x)^4]$ $= [1] - 8x + 24x^2 + [-32x^3 + 16x^4]$	M1		Any valid method as far as term(s) in x and term(s) in x^2 .
		A1		$p = -8$ Accept $-8x$ even within a series.
		A1	3	$q = 24$ Accept $24x^2$ even within a series.
(b)	x term is $\binom{9}{1} 2^8 x$ Coefficient of x term is $= 9 \times 2^8 = 2304 (=k)$	M1		OE
(c)	$(1-2x)^4 (2+x)^9 = (1+px+\dots)(2^9+kx\dots)$ $= \dots$ $= \dots + kx + px(2^9) + \dots$ Coefficient of x is $k + 512p$ $= 2304 - 4096 = -1792$	M1 M1	2	Condone $2304x$
		A1ft	3	Uses (a) and (b) oe (PI) Multiply the two expansions to get x terms ft on candidate's values of k and p . Condone $-1792x$ SC If 0/3 award B1ft for $p+k$ evaluated
	Total		8	

5(a)	$ar = 48; ar^3 = 3$ $\Rightarrow 16r^2 = 1$ $r^2 = \frac{1}{16} \Rightarrow r = -\frac{1}{4}$ or $r = \frac{1}{4}$	B1 M1 A1 B1		For either. OE Elimination of a OE CSO AG Full valid completion. SC Clear explicit verification (max B2 out of 3.)
(b)(i)	$a = -192$	B1	1	
(ii)	$\frac{a}{1-r} = \frac{a}{1-\left(-\frac{1}{4}\right)}$ $S_\infty = \frac{-768}{5} (= -153.6)$	M1 A1ft	2	$\frac{a}{1-r}$ used Ft on candidate's value for a . i.e. $\frac{4}{5}a$ SC candidate uses $r = 0.25$, gives $a = 192$ and sum to infinity = 256. (max. B0 M1A1)
	Total		7	

7(a)	$(1+2x)^8$ $= 1 + \binom{8}{1}(2x)^1 + \binom{8}{2}(2x)^2 + \binom{8}{3}(2x)^3 + \dots$ $= 1 + 16x + 112x^2 + 448x^3 + \dots$ $\{a = 16, b = 112, c = 448\}$	M1 A1A1 A1		Any valid method. PI by correct value for a, b or c A1 for each of a, b, c
(b)	x^3 terms <u>from expn.</u> of $\left(1+\frac{1}{2}x\right)(1+2x)^8$ $are \ cx^3 \ and \ \frac{1}{2}x(bx^2)$ $cx^3 + \frac{1}{2}x(bx^2)$	M1 A1		Either b, c or candidate's values for b and c from (a)
	Coefficient of x^3 is $c + 0.5b = 504$	A1ft	3	Ft on candidate's $(c + 0.5b)$ provided b and c are positive integers >1
Total			7	

2(a)	$u_1 = 12$ $u_2 = 3 \times 4^2 = 48$	B1 B1	2	CSO AG (be convinced)
(b)	$r = 4$	B1	1	
(c)(i)	$\{S_{12} = \frac{a(1-r^{12})}{1-r}$ $= \frac{12(1-4^{12})}{1-4}$ $= \frac{12(1-4^{12})}{-3} = -4(1-4^{12}) = 4^{13} - 4$	M1 A1ft A1	3	OE Using a correct formula with $n = 12$ Ft on answer for u_1 in (a) and r in (b) CAO Accept $k = 13$ for 4^{13} term
(ii)	$\sum_{n=2}^{12} u_n = (4^{13} - 4) - u_1$ $= 67108848$	B1	1	
Total			7	

4(a)	$\{S_{29} = \frac{29}{2}[2a + 28d]\}$	M1		Formula for S_n with $n = 29$ substituted and with a and d
	$29(a + 14d) = 1102$	m1		Equation formed then some manipulation
	$a + 14d = \frac{1102}{29} \Rightarrow a + 14d = 38$	A1	3	CSO AG
(b)	$u_2 = a + d$ $u_7 = a + 6d$	B1		Either expression correct
	$u_2 + u_7 = 13 \Rightarrow 2a + 7d = 13$	M1		Forming equation using u_2 & u_7 , both in form $a + kd$
	e.g. $21d = 63$; $3a = -12$	m1		Solving $a + 14d = 38$ with candidate's ' $2a + 7d = 13$ ' to at least stage of elimination of either a or d
	$a = -4$ $d = 3$	A1	4	Both correct
	Total		7	